

A relativistic quark model of baryons

Bernard Metsch, Ulrich Löring

*Institut für Theoretische Kernphysik,
Universität Bonn, Nußallee 14-16, D 53115 Bonn, Germany
E-mail: metsch@itkp.uni-bonn.de, loering@itkp.uni-bonn.de*

(Received: February 1, 2008)

On the basis of the three-particle Bethe-Salpeter equation we formulated a relativistic quark model for baryons. Assuming the propagators to be given by their free form with constituent quark masses and the interaction kernel by an instantaneous potential, which contains a string-like parameterization of confinement and a flavor dependent interaction motivated by instanton effects we can account for the major features in the baryon spectrum, such as the low position of the Roper resonance and the occurrence of approximate parity doublets apparent in the N - and Λ -spectra.

1 Introduction

Still the most successful description of the baryonic excitation spectrum is the non-relativistic constituent quark model (NRCQM). It essentially assumes that baryon resonances are q^3 -bound states, coupled weakly to meson fields. This certainly can be questioned: The decay widths of hadronic resonances are in general appreciable. The assumption should be understood to state that the coupling to strong decay channels does not influence the *relative* positions of resonances at the level of, say, 50 MeV. Thus the major mass splittings are supposed to reflect the excitation of the degrees of freedom of three constituent quarks, which have an effective constituent quark mass, carry flavor and color in the fundamental representations of the corresponding groups, are described by Pauli-spinors and thus obey the Pauli-principle. In its simplest form, assuming the interaction to be given by harmonic forces, these ingredients provides a natural explanation of the ground state flavor octet and decuplet, the lowest orbital excitations (“ $1-\hbar\omega$ -states”) and the rough position of the lowest state of each spin. A comparison with the experimental spectrum, however, also reveals some conspicuous shortcomings: there is no natural explanation of the position of the lowest scalar/isoscalar excitations, such as the Roper-resonance, nor of the low position of some radial/orbital Δ -resonances at 1.9 GeV. In addition to these phenomenological considerations there are also some more fundamental objections: even with constituent quark masses of a few hundred MeV, quarks in hadrons are not really slow—the expectation value of $\frac{p}{m_q}$ being roughly of order unity—processes at larger momentum transfer intrinsically require a relativistically covariant treatment of boosts, and in particular for the ground state mesons there are binding effects which are large compared to the masses of the constituents.

These considerations led us to the formulation of a relativistically covariant quark model for mesons on the basis of the instantaneous Bethe-Salpeter equation, see e.g. [1, 2] and references therein, while at the same time retaining all the successful features of the NRCQM.

2 The covariant q^3 Bethe Salpeter equation

In momentum space the Bethe-Salpeter equation for the Bethe-Salpeter amplitude $\chi_{\bar{P}}(x_1, x_2, x_3) = \langle 0 | T\Psi(x_1)\Psi(x_2)\Psi(x_3) | \bar{P} \rangle$ of a bound state with $\bar{P}^2 = M^2$ reads [3]

$$\begin{aligned} \chi_{\bar{P}}(p_\xi, p_\eta) = & S_1^F\left(\frac{1}{3}\bar{P} + p_\xi + \frac{1}{2}p_\eta\right) \otimes S_2^F\left(\frac{1}{3}\bar{P} - p_\xi + \frac{1}{2}p_\eta\right) \otimes S_3^F\left(\frac{1}{3}\bar{P} - p_\eta\right) \\ & \times (-i) \int \frac{d^4 p'_\xi}{(2\pi)^4} \frac{d^4 p'_\eta}{(2\pi)^4} K(\bar{P}, p_\xi, p_\eta, p'_\xi, p'_\eta) \chi_{\bar{P}}(p'_\xi, p'_\eta), \end{aligned} \quad (1)$$

where $\bar{P} = \sum_{i=1}^3 p_i$ is the total 4-momentum and $p_\xi = \frac{1}{2}(p_1 - p_2)$, $p_\eta = \frac{1}{3}(p_1 + p_2 - 2p_3)$ are Jacobi momenta. S^F denotes the full quark propagator and $K = K^{(3)} + \bar{K}^{(2)}$ is the integral kernel which contains both irreducible three-body $K^{(3)}$ and two-body forces $\bar{K}^{(2)}$. Both S^F and K are unknown functions for strongly interacting particles and thus have to be modeled. We start by making the assumption, that the full quark propagator can be approximated by its free form

$$S_i^F(p) \approx i [\gamma(p) - m_i + i\epsilon]^{-1},$$

where m_i is the effective constituent quark mass, which thus is a free parameter in our model. We furthermore assume, that the irreducible kernel can be suitably written as

$$K^{(3)}(p_\xi, p_\eta; p'_\xi, p'_\eta) |_{\bar{P}=(M, \vec{0})} = V^{(3)}(p_{\xi\perp}, p_{\eta\perp}; p'_{\xi\perp}, p'_{\eta\perp}), \quad K^{(2)}(p_\xi, p'_\xi) |_{\bar{P}=(M, \vec{0})} = V^{(2)}(p_{\xi\perp}, p'_{\xi\perp}) \quad (2)$$

where $p_\perp \equiv p - \frac{\bar{P} p}{\bar{P}^2} \bar{P}$. In a frame where $\bar{P} = (M, \vec{0})$ the kernel then depends on the spatial components of the relative momenta only and thus reflects an instantaneous potential in the rest frame of the baryon. Both assumptions are in fact borrowed from the NRCQM, which accounts for confinement by assuming a string-like three body potential for constituent quarks. In fact the instantaneous approach is somewhat problematic for two particle interactions. For the moment we will assume that retardation effects of the propagation of the spectator quark can be parametrized by construction of an effective 3-body kernel through $\langle G_0 \rangle V_{ij}^{\text{eff}} \langle G_0 \rangle := \langle G_0 K_{ij}^{(2)} G_0 \rangle$, where $G_0 := S_1^F \otimes S_2^F \otimes S_3^F$ is the free 3-quark propagator and $\langle G_0 \rangle := \int \frac{dp_\xi^0}{2\pi} \frac{dp_\eta^0}{2\pi} G_0$. With these assumptions one can integrate out the p_ξ^0, p_η^0 dependence in Eq.(1) and derive the Salpeter equation (see Ref. [3]):

$$\Phi = -i \langle G_0 \rangle (V^{(3)} + V_{\text{eff}}) \Phi \quad (3)$$

for the Salpeter amplitude $\Phi := \int \frac{dp_\xi^0}{2\pi} \frac{dp_\eta^0}{2\pi} \chi$, with $V_{\text{eff}} = \sum_{i < j} V_{ij}^{\text{eff}}$. The normalization of Φ follows from the normalization of the Bethe-Salpeter amplitude and reads

$$\sum_{\alpha\beta\gamma} \int \frac{d^3 p_\xi d^3 p_\eta}{(2\pi)^6} \Phi_{M\alpha\beta\gamma}^*(\vec{p}_\xi, \vec{p}_\eta) \Phi_{M\alpha\beta\gamma}(\vec{p}_\xi, \vec{p}_\eta) = 2M. \quad (4)$$

This allows to reformulate Eq.(3) as an eigenvalue problem $\mathcal{H} \Phi = M \Phi$ with the Salpeter Hamiltonian

$$(\mathcal{H} \Phi_M)(\vec{p}_\xi, \vec{p}_\eta) = \sum_{i=1}^3 H_i(\vec{p}_i) \Phi_M(\vec{p}_\xi, \vec{p}_\eta) + [\Lambda_1^+(\vec{p}_1) \otimes \Lambda_2^+(\vec{p}_2) \otimes \Lambda_3^+(\vec{p}_3) + \Lambda_1^-(\vec{p}_1) \otimes \Lambda_2^-(\vec{p}_2) \otimes \Lambda_3^-(\vec{p}_3)] \gamma^0 \otimes \gamma^0 \otimes \gamma^0 \int \frac{d^3 p'_\xi}{(2\pi)^3} \frac{d^3 p'_\eta}{(2\pi)^3} (V^{(3)} + V_{\text{eff}})(\vec{p}_\xi, \vec{p}_\eta, \vec{p}'_\xi, \vec{p}'_\eta) \Phi_M(\vec{p}'_\xi, \vec{p}'_\eta) \quad (5)$$

with the Dirac Hamiltonian H_i and $\Lambda_i^\pm$ being projectors on states of positive (+) and negative (−) energy spinors. Note that with the present instantaneous approximation we get the same number of states as in the NRCQM, nevertheless our approach goes beyond a simple "relativization" by explicitly taking into account the coupling to negative energy components. In this formulation we can determine masses and Salpeter amplitudes for baryons by diagonalizing the Hamiltonian of Eq.(5) in an appropriate large, but finite basis. In order to calculate electroweak observables in the framework of the Mandelstam formalism we need the full Bethe-Salpeter amplitude. This can be found in the rest frame through the Bethe Salpeter equation (see Ref. [11])

$$\chi_M = -i G_0 (V_{\text{conf}}^{(3)} + V^{\text{eff}}) \Phi,$$

and, because of formal covariance, can be calculated for any on-shell momentum $\bar{P}$ with $\bar{P}^2 = M^2$.

3 A covariant quark model for light-flavored baryons

In order to calculate the spectrum of baryons with light flavors we have to specify the interaction kernels $K^{(3)} = V^{(3)}$ and $K^{(2)} = V^{(2)}$ in instantaneous approximation. We parameterize a three-body confinement kernel by a potential $V_{\text{conf}}^{(3)}$ which rises linearly with inter-quark distances. The gross features of the baryon resonances seem to indicate that the dominating confinement forces should be spin-independent, at least in the non-relativistic limit; moreover, too large spin-orbit effects should be suppressed. To realize both properties we assume the Dirac structure given by a specific combination of spin-independent scalar and time-like vector¹ Dirac structures:

$$V_{\text{conf}}^{(3)}(\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3) = 3a \frac{1}{4} [\mathbb{1} \otimes \mathbb{1} \otimes \mathbb{1} + \gamma^0 \otimes \gamma^0 \otimes \mathbb{1} + \text{cycl. perm.}] + b \sum_{i < j} |\mathbf{x}_i - \mathbf{x}_j| \frac{1}{2} [-\mathbb{1} \otimes \mathbb{1} \otimes \mathbb{1} + \gamma^0 \otimes \gamma^0 \otimes \mathbb{1} + \text{cycl. perm.}] . \quad (6)$$

Here, the offset a and the slope b enter as free parameters in our model.

In order to describe spin-dependent splittings, such as that of the ground-state octet and decuplet, we adopt the explicitly flavor-dependent 2-quark interaction², derived by 't Hooft from instanton effects:

$$V_{\text{'t Hooft}}^{(2)}(\mathbf{x}) = \frac{-4}{\lambda^3 \pi^{\frac{3}{2}}} e^{-\frac{|\mathbf{x}|^2}{\lambda^2}} [\mathbb{1} \otimes \mathbb{1} + \gamma^5 \otimes \gamma^5] \mathcal{P}_{S_{12}=0}^{\mathcal{D}} \otimes (g_{nn} \mathcal{P}_{\mathcal{A}}^{\mathcal{F}}(nn) + g_{ns} \mathcal{P}_{\mathcal{A}}^{\mathcal{F}}(ns)) . \quad (7)$$

Here, the four-fermion contact interaction has been smeared out by a Gaussian function with effective range λ ; the operator $\mathcal{P}_{S_{12}=0}^{\mathcal{D}}$ in Dirac space projects on anti-symmetric spin-singlet states, $\mathcal{P}_{\mathcal{A}}^{\mathcal{F}}(nn)$ and $\mathcal{P}_{\mathcal{A}}^{\mathcal{F}}(ns)$ denote the projectors on flavor-antisymmetric non-strange and non-strange-strange quark pairs. Although the constituent quark masses m_n, m_s as well as the 't Hooft couplings g_{nn}, g_{ns} and the range λ can be related to standard QCD-parameters by instanton theory, we determine these parameters by a fit to the experimental baryon spectrum; for the values of the seven free parameters and a consistency check of the QCD-relations we refer to Refs. [4, 5].

As 't Hooft's interaction affects flavor antisymmetric qq-pairs only this interaction does not act on flavor symmetric states, such as the Δ -resonances, which are thus determined by the dynamics of the confinement potential alone. Accordingly, the constituent quark masses and the confinement parameters were determined by a fit to the spectrum in this sector without any influence of the residual interaction. In this manner a reasonable description of the complete Δ -spectrum including the linear Regge trajectory ($M^2 \sim J$) up to highest orbital excitations $J = \frac{15}{2}$ can be obtained, see the upper part of Fig. 1. However, the suspicious low position of the three negative-parity “ $3\hbar\omega$ ” states quoted by the PDG [6] around 1900 MeV as well as that of the $\Delta_{\frac{3}{2}}^{+}(1600)$ is not accounted for (as in several other quark models). Since in our opinion the experimental evidence of these negative parity states from π - N -scattering is rather weak, a confirmation by complementary (electro-/photo-production) experiments, such as the CLAS experiment at CEBAF (JLab) or the Crystal Barrel experiment at ELSA (Bonn), seems of great importance.

The instanton-induced interaction does act on particular flavor octet (and singlet) states. The 't Hooft coupling g_{nn} is adjusted to the ground-state N - Δ -splitting (and correspondingly g_{ns} to the

¹The appearance of $\gamma^0 \otimes \gamma^0$ is allowed because the potential is defined in the rest-frame of the baryon.

²This interaction is usually used in connection with the solution of the $U_A(1)$ -problem. In the framework of calculations for mesons [1] we have indeed shown that it yields the correct splitting of the lowest meson nonet and in general of all low-lying meson states. Moreover, we found [4] that the alternative QCD-based residual interaction – the one-gluon exchange – can be discarded on phenomenological grounds.




Figure 1: The complete Δ^- (above) and nucleon spectrum (below): Our calculation in the left part of each column is compared to the experimental spectrum [6] in the right part of each column. Resonances are classified by total spin and parity J^π . The experimental position is indicated by a bar, the corresponding uncertainty by the shaded box which is darker for better established resonances; the status of each resonance is additionally indicated by stars.

$\Xi^*-\Xi^-$ and $\Sigma^*-\Sigma-\Lambda$ -splittings in the strange sector); then all other states of the nucleon- (and the Λ^- , Σ^- , Ξ^-) spectrum are real parameter-free predictions. The lower part of Fig. 1 shows as an example our predictions for the complete N -spectrum. As can be seen from this figure, our predictions can nicely reproduce several features at least in qualitative but mostly even in completely quantitative agreement with the experimental findings.

It is instructive to analyze the consequences of instanton effects for the shape of the spectrum in some more detail. Figure 2 illustrates the effect of 't Hooft's force on the lowest nucleon states. We find a selective lowering of some particular states and once the coupling is fixed to reproduce the experimental N -ground-state position we can automatically account very well for the major spin-dependent mass splittings observed in the experimental spectrum. For instance in the " $2\hbar\omega$ " band exactly four states are lowered relative to the other states such as required by the experimental findings. In particular one finds that in this manner the extremely low position of the prominent

Figure 2: Instanton-induced effects in the nucleon spectrum. The curves illustrate the change of the spectrum with increasing 't Hooft coupling g_{nn} which finally is fixed to reproduce the correct N - Δ -splitting.





Figure 3: Parity doublets for higher lying nucleon resonances. See also caption to Fig. 2.



Roper resonance can be accounted for quite naturally. We should remark here that the specific interplay of 't Hooft's force, the relativistic effects and the particular choice for the Dirac structure of the confinement potential are essential to describe these effects quantitatively [4].

A very interesting feature of our relativistic quark model with instanton induced forces is that it gives a natural dynamical explanation of the so-called *parity doublets*: A glance at the experimental N - and Λ -spectrum reveals a conspicuous degeneracy of some states with the same spin and opposite parity. Prominent, well-established, examples are $N_{\frac{5}{2}}^{5+}(1680)$ - $N_{\frac{5}{2}}^{5-}(1675)$, $N_{\frac{9}{2}}^{9+}(2220)$ - $N_{\frac{9}{2}}^{9-}(2250)$, $\Lambda_{\frac{5}{2}}^{5+}(1820)$ - $\Lambda_{\frac{5}{2}}^{5-}(1830)$. In the Σ -spectrum no clear indications of parity doublets is found. In the literature these observations have been related to a phase transition from the Nambu-Goldstone mode of chiral symmetry to the Wigner-Weyl mode in the upper part of the baryon spectrum [7-9]. In the present model also this feature can be understood as an instanton-induced effect: In general 't Hooft's force lowers those sub-states of a major oscillator shell³ which contain so-called scalar diquarks, i.e. quark pairs with trivial spin and angular momentum. This is found in particular for the high spin states in a given $N\hbar\omega$ oscillator shell, see Fig. 3. For given $N\hbar\omega$ the maximum total angular momentum for a state containing such a scalar diquark is $J = (L_{\max} = N) + \frac{1}{2}$. 't Hooft's force lowers this state enough to become almost degenerate with the unaffected spin-quartet state of the adjacent oscillator shell with $N - 1$, which has opposite parity but the same total angular momentum: $J = (L_{\max} = N - 1) + \frac{3}{2}$. In this way patterns of approximate parity doublets for *all* lowest excitations in the sectors $J = \frac{5}{2}$ to $J = \frac{13}{2}$ are formed *systematically*. In the $N_{\frac{5}{2}}^{\pm}$

³Note that despite of the linear confinement adopted here, the assignment of states to oscillator shells still provides an adequate classification of states with confinement alone.

and $N_{\frac{1}{2}}^{9\pm}$ sectors this scenario is nicely confirmed experimentally by the well-established parity doublets mentioned above: $N_{\frac{1}{2}}^{5+}(1680)$ – $N_{\frac{1}{2}}^{5-}(1675)$ and $N_{\frac{1}{2}}^{9+}(2220)$ – $N_{\frac{1}{2}}^{9-}(2250)$. In the $N_{\frac{1}{2}}^7$ sector, however, the present experimental findings seem to deviate from such a parity doubling structure due to the rather high position of the $N_{\frac{1}{2}}^7(2190)$. Although this state is given a four-star rating [6] an investigation of this sector with the new experimental facilities would be highly desirable. The same mechanism explains approximate parity doublet structures also for states with lower angular momentum as *e.g.* the N^* doublets in the second resonance region around ~ 1700 MeV with spins $J^\pi = \frac{1}{2}^\pm, \frac{3}{2}^\pm$, and $\frac{5}{2}^\pm$ (see Fig. 2). Observable consequences of this parity doubling scenario should manifest in a different shape of electromagnetic $\gamma^*p \rightarrow N^*$ transition form factors of both members of a doublet due to their significantly different internal structures: That member of the doublet, which is affected by 't Hooft's force (e.g. $N_{\frac{1}{2}}^{5+}$), exhibits a rather strong scalar diquark correlation and thus its structure should be more compact compared to its unaffected doublet partner (e.g. $N_{\frac{1}{2}}^{5-}$) whose structure is expected to be rather soft. Consequently, the transition form factor to the latter resonance decreases faster than that to its doublet partner with the scalar diquark contribution (see Fig. 3 of Ref. [10] for an example). Finally, we should mention that in the strange sector (see Ref. [5]) 't Hooft's force accounts in a similar way for the prominent doublets of the Λ -spectrum. At the same time instanton-induced effects are found to be significantly weaker in the Σ -spectrum, thus explaining the fact that no clear experimental indications of parity doublets are observed in this sector.

The investigations presented here have been restricted to the spectra of baryons alone, which poses of course only a first (successful) test of our model. Strong and electroweak decays as well as formfactors pose additional stringent tests. In the present covariant framework formfactors were already computed in the Mandelstam formalism [11]; indeed we found that we can reliably calculate such observables up to high momentum transfers. Work on the perturbative calculation of strong decays is more difficult but in progress; first results are very promising, e.g. a *parameter-free* (!) prediction [12] of $\Delta_{\frac{3}{2}}^+(1232) \rightarrow N\pi$ yields a decay width $\Gamma = 109$ MeV in very good agreement with the experimental value $\Gamma = 119 \pm 5$ MeV.

Acknowledgments: We would like to acknowledge the contributions of K. Kretzschmar, D. Merten and H.-R. Petry. We thank the Deutsche Forschungsgemeinschaft (DFG) for financial support.

References

- [1] M. Koll, R. Ricken, D. Merten, B.C. Metsch, H.R. Petry, Eur. Phys. J. A **9**, 73 (2000).
- [2] R. Ricken, M. Koll, D. Merten, B.C. Metsch, H.R. Petry, Eur. Phys. J. A **9**, 221 (2000).
- [3] U. Löring, K. Kretzschmar, B.Ch. Metsch, H.R. Petry, Eur. Phys. J. A **10**, 309 (2001).
- [4] U. Löring, B.Ch. Metsch, H.R. Petry, Eur. Phys. J. A **10**, 395 (2001).
- [5] U. Löring, B.Ch. Metsch, H.R. Petry, Eur. Phys. J. A **10**, 447 (2001).
- [6] Particle Data Group, 'Review of Particle Physics', Eur. Phys. J. **C15**, 1-878 (2000).
- [7] L. Y. Glozman, Phys. Lett. B **475**, 329 (2000).
- [8] T. D. Cohen and L. Y. Glozman, hep-ph/0102206 (2001).
- [9] M. Kirchbach, Nucl. Phys. A **689**, 157 (2001).
- [10] U. Löring, B.Ch. Metsch, *Parity doublets from a relativistic quark model*, Proceedings of the *Workshop on The Physics of Excited Nucleons (NSTAR 2001)*, World Scientific (2001), p.221.; see also hep-ph/0110412
- [11] K. Kretzschmar, U. Löring, B.Ch. Metsch, *Elektroweak form factors of the nucleon in a relativistic quark model of baryons*, in progress.
- [12] D. Merten, private communication (2001)